

# Capacity Charts

## As-fed upright silo capacity

Size, ft (diameter x height)	Corn silage & haylage			HMSC	Ground HMSC	Ground HM ear corn
Moisture content	70%	60%	50%	30%	30%	30%
12x30	80	65	50	89	95	70
12x40	115	90	70	120	128	94
12x50	155	120	95	151	160	120
14x40	160	120	96	165	172	128
14x50	213	160	127	208	220	163
14x60	266	200	160	251	264	198
16x30	146	110	88	150	166	123
16x40	209	156	125	220	224	167
16x50	278	208	167	274	285	213
16x60	347	260	208	330	345	259
18x40	265	198	159	270	284	211
18x50	352	264	211	350	360	269
18x60	440	330	264	422	437	328
18x70	530	398	318	496	520	389
20x40	326	245	196	340	350	260
20x50	435	326	261	428	445	332
20x60	543	407	326	525	540	404
20x70	655	491	393	617	638	480
20x80	767	575	460	708	736	557
24x50	626	470	375	600	640	478
24x60	782	587	469	763	776	582
24x70	943	707	565	897	918	692
24x80	1,104	828	662	1,032	1,060	801
24x90	1,275	955	764	1,165	1,209	920
30x80	1,725	1,293	1,035	1,628	1,656	1,252
30x90	1,990	1,493	1,195	1,840	1,888	1,434

## Wagon capacity

Depth, ft	Approximate tons (as-is basis)							
	Length, ft (65% moisture)				Length, ft (55% moisture)			
	14	16	18	20	14	16	18	20
3	3.5	4.0	4.5	5.0	2.5	3.0	3.5	4.0
4	4.5	5.5	6.0	6.5	3.5	4.0	4.5	5.0
5	6.0	6.5	7.5	8.5	4.5	5.0	5.5	6.5
6	7.0	8.0	9.0	10.0	5.5	6.0	7.0	7.5
7	8.0	9.5	10.5	12.0	6.0	7.0	8.0	9.0
8	9.5	11.0	12.0	13.5	7.0	8.0	9.0	10.0

## Bag capacity\*

Bag size, ft	Tons (fresh)
8	1
9	1.25
10	1.5
11	1.75
12	2.25
14	2.75

\*Estimates shown are for corn silage.

## Ratio for bag chart

Type of crop, moisture	Relation to 65% corn silage tonnage
Haylage, 60%	100%
HMSC, 30%	130%
Earlage, 35%	120%
Snaplage, 40%	130%

## Estimated as-fed capacity for bunkers and piles

Enter avg width*	1. _____	ft
Enter avg length	2. _____	ft
Enter avg height	3. _____	ft
Multiply 1x2x3	4. _____	lb/ft <sup>3</sup> in structure
Est as-fed density**	5. _____	lb/ft <sup>3</sup>
Multiply 4x5	6. _____	lb as-fed in structure
Divide by 2,000	7. _____	tons as-fed in structure

\*To determine dimensions for piles, look at the slopes of each side of the pile. Visualize how much of the slope would need to be "folded back" on itself to square up the sides of the pile to determine average width.

\*\*Use known as-fed density when possible. Otherwise, start with these average densities: 40 lb for haylage and corn silage; 60 lb for HMSC; and 45 lb for earlage/snaplage. Use higher or lower numbers for well packed or poorly packed units, respectively.

# CROP-N-RICH<sup>®</sup> NEXUS

Easy-to-use dual-purpose  
silage inoculant for greater  
dry matter recovery and  
reduced spoilage and heating  
in a broad range of forages

Contains  
*Lactobacillus plantarum* MTD/1<sup>®</sup>  
and *Lactobacillus buchneri* PJB/1



Service & technology to get  
the most from your forage

[www.vitaplus.com/forage-foundations](http://www.vitaplus.com/forage-foundations)



Vita Plus Corporation  
PO Box 259126, Madison, WI 53725-9126  
800.362.8334 • [www.vitaplus.com](http://www.vitaplus.com)

REV 0823





# CROP-N-RICH® NEXUS

Dual-purpose silage inoculant with *L. plantarum* MTD/1® and *L. buchneri* PJB/1 bacteria improves dry matter (DM) recovery and reduces spoilage and heating

## Proven effective

### Research shows:

- Improved upfront fermentation and DM recovery
- Less protein degradation and ammonia production
- Less spoilage, longer bunklife and less wasted feed
- Improved palatability

### Effective dual-purpose bacterial inoculant:

1. ***L. plantarum* MTD/1** is the most research-proven silage inoculant bacteria that produces lactic acid to quickly lower the silage pH and reduce DM loss.
  - Crop-N-Rich Nexus supplies 100,000 colony forming units (CFU) of *L. plantarum* MTD/1 per gram of forage.
2. Acetic acid is a powerful fungicide that reduces yeast growth and subsequent mold, spoilage, and heating. ***L. buchneri* PJB/1** works by producing higher levels of acetic acid in the silage.
  - Crop-N-Rich Nexus supplies 200,000 CFU of *L. buchneri* PJB/1 per gram of forage.

## Unmatched ease of use

- Water-soluble prills easily mix into solution in less than 45 seconds


### Use when:

- Forages are fed during warm weather, feedout rates are a challenge, high-starch feeds are stored, feeds may be moved after storage, TMRs tend to heat in the feedbunk, or crops have been compromised by rain, hail, insects, drought, or disease
- Baleage or haylage is stored with greater than 35% DM and corn silage is stored with greater than 32% DM
- Feedout will begin after at least 45 days in storage to maximize aerobic stability benefits

### Product options:


- **Crop-N-Rich Nexus 100:** Treats 100 tons (as-is) of forage
- **Crop-N-Rich Nexus 500:** Treats 500 tons (as-is) of forage

*Lower pH and improved DM recovery in grass (42% DM) treated with Crop-N-Rich Nexus.*



Source: Internal trial data.

*Fewer yeasts and improved aerobic stability in corn silage (32% DM) treated with Crop-N-Rich Nexus at 30 and 90 days post-ensiling.*



Source: University of Delaware, 2014.

*Average of 32 trials with corn silage, wheatlage, alfalfa, and earlage harvested at DM ranging from 28% to 62% and treated with Crop-N-Rich Nexus.*

	Control	Crop-N-Rich Nexus
Acetic acid, %	1.42	2.04
Yeasts, CFU/g	39,346,562	149,234
Aerobic stability, hours	104	236